Glycolysis

Glycolysis ÷ — 1-5 : Energy-investment
— 6-10 : Energy-generation (Sub.-level)

Net products: 1 Glucose → 2 Pyruvate
2 ADP → 2 ATP
2 NAD^+ → 2 NADH

Amphibol-P : 2 + 4 = 6
Malate-Aspartate : 2 + 6 = 8

Glycogen ⇌ Glucose (Glycogenesis / Glycogenolysis) ⇌ Pyruvate (Glycolysis / Gluconeogenesis)

GLUT TRANSPORTERS

I- BBB — B- Blood (RBCs - Erythrocytes)
B- Baby (Tissue)
B- Blood-brain barrier

II- KIDS LIPS — K- Kidney, L- Liver, P- Pancreas

III- PINE (or PINK) — P - Placenta, N - Neuron, E - Kidney

} insulin-independent

IV- MOTHER — M- Muscle (Skeletal muscle)
FATHER — F- Fat (Adipose tissue)

} insulin-dependent

Glycolysis:
- Occurs in cytoplasm
- S.S: Glucose
- E.P: 2 Pyruvates
- Net ATP: 2 ATP
- 2 NADH

Glucose
(Liver) glucokinase / hexokinase (muscles & other tissues)
ATP → ADP
Glucose-6-P
phosphoglucose isomerase
Fructose-6-P
phosphofructokinase I (rate-limiting & committed step of glycolysis)
ATP → ADP
Fructose-1,6-BP
aldolase
DHAP ⇌ (triose-P isomerase) Glyceraldehyde-3-P
2 NAD^+ + 2 H_3PO_4 → 2 NADH
1,3-bisphosphoglycerate
2 ADP + 2 ATP — phosphoglycerate kinase
3-phosphoglycerate
phosphoglycerate mutase
2-Phosphoglycerate
enolase
Phosphoenol Pyruvate
ADP + ATP — pyruvate kinase
Pyruvate

Glucose → (glucokinase/hexokinase) Glucose-6-P → (phosphohexose isomerase) Fructose-6-P → (6-phosphofructokinase I (6-PFK-1)) Fructose-1,6-BP → (aldolase) DHAP + Glyceraldehyde-3-P → (triose phosphate isomerase) 1,3-Bisphosphoglycerate → (phosphoglycerate kinase) 3-Phosphoglycerate → (phosphoglycerate mutase) 2-Phosphoglycerate → (enolase) PEP → (Pyruvate kinase) Pyruvate

pyruvate kinase deficiency → hemolytic anemia

Regulatory key enzymes: (3)

1) Glucokinase / hexokinase : Glucose → Glucose-6-Phosphate (+ATP / +ADP+Pi)

2) 6-Phosphofructokinase-1 : Fructose-6-Phosphate (+ ATP) → fructose-1,6-BP (+ ADP+Pi)
• MAIN REGULATING

3) Pyruvate Kinase : PEP → Pyruvate (ADP → ATP)

Activation

key / Glu :
- insulin
- Fru-1-P

- ↓ ATP/AMP
- Fru-2,6-BP
- insulin

- insulin
- Fru-1,6-BP

Inhibition

Glu-6-P
Fru-6-P

- ↑ ATP/AMP
- citrate
- acidic pH

- Glucagon
- ↑ ATP/AMP
- acetyl-CoA

Pyruvate → PEP ⇌ 2-phospho-glycerate ⇌ 3-phospho-glycerate ⇌ 1,3-BP glycerate ⇌ Interchangeable 3-P ⇌ fructose-1,6-BP

in DHAP

↓

fructose-6-phosphate

↓

glucose-6-phosphate

↓

glucose

Gluconeogenesis

" reversed glycolysis "

w/ exception of 3 reactions

Breakdown of lipids, ketone bodies

Energetically important lipids → TAGs
(while not discussing phospholipids ...)

ox. of 1g of { fatty acids = 38 kJ / carbs. / proteins } 17 kJ

TAG supply sufficient for ~week

healthy person has 15% of body weight = TAGs

MAIN SITE OF TAG ACCUMULATION = adipocytes (fat cells)

Lipid breakdown:

Lipids as E source

hormone sensitive lipase (HSL) → release FAs from store,
performs complete lipolysis
1 TAG → 3 FA + 1 glycerol

inhibited by: insulin
activated by: counter-reg. hormones →
glucagon, adrenaline, noradrenaline

insulin wants less fats & stuff →
doesn't want to release
FAs & glycerol
& opposite for glucagon, adrenaline, noradrenaline

utilizing lipids → 3 steps

1) lipid mobilization : TAG → DFA + glycerol transported by blood
2) activation of FA : transp. from cytosol (where they're activated) to MIT matrix
3) β-oxidation : FA → acetyl-CoA → CO_2
reduced cofactors produced, used dietary in resp. chain

1) lipid mobilization

TAG → DFA + glycerol
released from fat into blood

FA longer than 12C →
cannot in plasma evenly.
(hydrophobic)
(albumin → albumin)

glycerol → in liver (glycolysis / gluconeogenesis)
glycerol → glycerol-3-P → DHAP
ATP, kinase; glycerol-3-P DH

Lipolysis (regulation)	Activation	Inhibition
hormone sensitive lipase (adipocytes)	· (adrenaline/cortisol) · glucagon · ACTH (phosphorylation)	· insulin · prostaglandins
lipoprotein lipase (endothelium surface)	· insulin · apoliproteins C-II	

Metabolism of ketone bodies

3 ketone compounds

acetoacetate, β-hydroxybutyrate, acetone

(acetoacetate, β-hydroxybutyrate: essential; acetone: waste product, excreted)


- completely degraded to CO_2 & water

Ketones essentially blood-transportable version of acetyl (they're derived from acetyl-CoA)

ONLY formed in liver mitochondria

so ketogenesis (ketone body synthesis) → organ: liver ONLY → cellular location: mitochondria ONLY

ketone bodies:



glucose runs out (starvation) → ketones produced by liver

ketones used by all body except liver (no enzyme)

& erythrocytes (RBCs) → no mito. → no ketone body breakdown (degradation)

carboxylase deficiency → OAA deficiency → TCC slows down → more convert acetyl-CoA → ketogenesis

& low glycemia → glucagon activated → TAG breakdown bec. HSL activated by glucagon → high FA → β-oxidation → high acetyl-CoA → ketogenesis

Energy Storage in cells

After meal → ↑ insulin/glucagon ratio → ↑ nutrients converted to ↑ storage molecules

Excess of glucose → ↑ stored as glycogen
→ ↑ converted to pyruvate (glycolysis) then acetyl-CoA (used for FA synthesis).

Excess of FA (from diet or formed from excess glucose & AA) → ↑ synthesized into TAGs (main energy reserve form in body)

fat storage → TAG
saccharide storage → ↑ glycogen
protein storage → ↑ none

Glycogen → Glucose storage in animal cytoplasm (violet color) in almost all cells

women → ↑ more glycogen → ↑ vaginal mucosa

Glycogen in: liver → used to maintain glycemia; muscle → used during short-term strenuous muscle work; vaginal mucosa → ↑ defense against microorganisms → ↑ acidic environment (glycogen → ↑ lactate → ↓ pH + protection).

PAS → ↑ stains glycogen brightly, violet.

Glycogen:
branched homopolymer
α-1,4-glycosidic bonds
branching: α-1,6 glycosidic bonds
2 groups of enzymes to cleave α1,4 & α1,6

glycogen → 2 ends
elongation & processing → ↑ non-reducing end (w/o OH)


reducing end was semi-acetyl mycrarg) bound by protein glycogenin

Glycogen: branched on every 8th - 10th glucose
only 1 reducing end, all left non-reducing (branches)

Glycogenesis → ↑ glycogen synthesis
- in cytosol
- 4 steps

1) phosphorylation: in liver: glucokinase; in muscle: hexokinase — glucose + ATP → Glc-6-P + ADP
2) isomerization: glucose phosphate isomerase — Glc-6-P → Glc-1-P (= coi ester)
3) coupling of UDP: UDP-glucose pyrophosphorylase — Glc-1-P + UTP → UDP-Glc (anabolic compound formed)
4) connection of UDP-Glc to non-reducing end of glycogen: glycogen synthase
UDP released, α-1,4 bond formed
8- to glucoses then branch

branching enzyme: branching enzyme = amylo-(1,4→1,6)-transferase



Glycogen synthesis:
- primer required: remaining glycogen or glycogenin enzyme

Regulation of glycogen synthesis: glycogenesis:

glycogen Synthase:
[active in dephosphorylated form
inactive in phosphorylated form

to phosphorylate glycogen synthase (inactive) → ↑ protein kinase
to dephosphorylate glycogen synthase (active) → ↑ phosphoprotein phosphatase

- insulin → wants to ↓ free glucose levels in blood → activates glycogen synthase
glucagon inhibits it
adrenaline inhibits it

protein kinase A (PKA) → ↑ cAMP dependent
(CAMK (calmodulin dependent protein kinase) - active when protein calmodulin binds)

4 Ca^{2+} → ↑ binds to calmodulin → ↑ active calmodulin can now bind to CAMK

glucagon → ↑ activates adenylate cyclase → ↑ forms cAMP → ↑ activates PKA → ↑ phosphorylate glycogen synthase
muscle contraction → ↑ (Ca^{2+}) intracellularly → ↑ binds to calmodulin → ↑ active calmodulin → ↑ activate/inactivate (CAMK) → ↑ inactivate) it
→ ↑ activates PKA → ↑ phosp. glyc. syn. → ↑ inactivates it

insulin → activates phosphodiesterase → cleaves cAMP to AMP → inactivates PKA

↳ also activates, phosphoprotein phosphatases → dephosp. → activates glyc synth.

Glycogenolysis (hydrolysis): degradation of glycogen molecule (breakdown)

- also in cytosol
- 3 steps

1) phosphorolytic cleavage

α1,4 bond cleaved → Glc-1-P (Cori ester) formed

free P$_i$ used linked to C$_1$ glucose

Enzyme: glycogen phosphorylase

ends 4 glc molecs. before branching

2) isomerization:

Glc-1-P ⟶ Glc-6-P
(phosphoglucomutase / glucose phosphate isomerase)

3) cleavage of branch → removal of α-1,6 bond

glucanotransferase (+ debranching enzyme)

Regulation of glycogen degradation (glycogenolysis):

glycogen phosphorylase → active: phosphorylated by phosphorylase kinase
→ inactive: dephosphorylated

phosphorylase kinase: inhibited by insulin, activated by counter regulatory hormones


Signaling via PKA & CAMK

in food (meal) → glycogen present → free glucose absorbed by blood (α-amylase (hydrolase))

Regulation	Activators	Inhibitors
Synthesis: Glycogen synthase	: Insulin, Glc-6-P	: calcium, Adrenaline
Degradation: Glycogen phosphorylase	• Glucagon, Adrenaline, ↓ATP, AMP, • Ca^{2+} (in muscle)	• ↑ATP, AMP • Glc-6-P • Glucose


Synthesis of fatty acids

FA degradation vs. synthesis



- FAD & NAD^+ e⁻ acceptors
- CoA is acyl group carrier

Biosynthesis - O in cytoplasm



- NADPH is e⁻ donor
- ACP is acyl carrier

Malonyl-CoA formation:

- it's the initial substrate for FA-acetyl-CoA synthesis

acetyl-CoA + ATP + HCO_3 → malonyl-CoA + ADP + Pi + H^+

carboxylation
enzyme: acetyl-CoA carboxylase
cofactor: biotin (vit. H1B7)

acetyl-CoA carboxylase → reg. enzyme of whole pathway → also affects FA β-oxidation

CO_2 removed during condensation w/ growing FA

Individual steps of FA synthesis:
(uses NADPH from pentose cycle)

- 1) malonyl-CoA synthesis w/ enzyme acetyl-CoA carboxylase

acetyl-CoA + ATP + HCO_3^- → malonyl-CoA + ADP + Pi + H^+

- 2) binding of acetyl to cond. enzyme (CE) w/ enzyme acetyltransacylase

acetyl-CoA + CE → acetyl-CE + CoA

- 3) malonyl-CoA binding to ACP (acyl carrier protein) w/ enzyme malonyltransacylase

malonyl-CoA + ACP → malonyl-ACP + CoA

- 4) condensation reaction: connecting malonyl ACP & acetyl-CE w/ enzyme β(3)-ketoacyl-ACP synthase

acetyl-CE + malonyl-ACP → acetoacetyl-ACP + CE + CO_2

- 5) reduction w/ enzyme β-ketoacyl reductase

acetoacetyl-ACP + NADPH → 3-hydroxybutyryl-ACP + H_2O

- 6) dehydration w/ enzyme 3-hydroxyacyl dehydratase

D-3-hydroxybutyryl-ACP → crotonyl-ACP + H_2O

- 7) reduction w/ enzyme enoyl reductase

crotonyl-ACP + NADPH → butyryl-ACP + NADP

end product: palmitate (16c)

thioesterase cleaves ACP from palmitate by hydrolyzing thioester bond

requirements for palmitate formation

8 acetyl-CoA
14 NADPH
7 ATP

acetyl-CoA transp. to mito using citrate

8 NADPH obtained by transp. citrate to cyto. & remaining 6 in pentose cycle

TAG synthesis:

TAG - final form of FA storage

- starting w/ DHAP

  reduction: DHAP OH

  DHAP + NADH → glycerol-3-P + NAD^+

- adding first 2 FAs

  enzyme: acyltransferase

  glycerol-3-P + acyl-CoA → monoacylglycerol-3-P + CoA

  monoacylglycerol-3-P + acyl-CoA → diacylglycerol phosphate (phosphatidic acid)

- remove P_i from phosphatidic acid hydrolytically

  enzyme: phosphatidic acid phosphatase

  phosphatidic acid + H_2O → 1,2-diacylglycerol + P_i

- connection of last FA

  enzyme: acyltransferase

  1,2-diacylglycerol + acyl-CoA → TAG + CoA

Metabolism of amino acids


Proteinogenic AAs:

biosynthesis: Essential - not made in body, get from diet
Non essential - made in body

degradation: Glucogenic: C skeleton makes Glc
Ketogenic: C skeleton makes Acetyl-CoA

Essential AAs: Val, Leu, Ile
PVT TIM HALL — Phe, (Tyr), Trp
His, Lys, Arg
Met (Cys)
Thr

Other important AAs in body:



AAs used for:
proteosynthesis
N-compound synthesis
degradation (E, glucose, fat)

AA degradation: 1) $-NH_2$ group removal
2) deaminization of amino group
3) C skeleton metabolism
→ 7 products

cofactor for amino transferases:
pyridoxal phosphate (PLP) derived from vit. B_6 (pyridoxine)

transamination:
$AA_1 + \alpha\text{-}KA_2 \rightarrow \alpha\text{-}KA_1 + AA_2$

alanine + glutamate ⟷ α-KG + pyruvate

OAA
Asp
AST or GOT
alanine aminotransferase (ALT) or GPT
α-KG

Metabolism of lipoproteins

Absorbed fat (diet) & lipids synthesized in liver & adipose tissue → must be transported between various tissues & organs to use & store.

Lipids → insoluble in water
associate (TAGs & CE) w/ amphipathic lipids (phospholipids & cholesterol) & proteins to make water-miscible lipoproteins
nonpolar lipids

Lipoprotein = lipid + apoprotein

Cholesterol → Good - HDL
Bad - LDL

Lipoproteins classified based on density

Chylomicrons
VLDL
IDL
LDL
HDL
FFA / NEFA

Lipoproteins are classified according to proportion of individual components that affect their density

TAGs - lowest density
proteins - highest density

Surface of lipoproteins has monolayer of phospholipids w/ free cholesterol and Apolipoproteins (Apo)
inside: TAG & CE

Chylomicrons:
lowest density
most TAG, least proteins

Chylomicrons
B-48
C-II
E
produced in intestine
principal lipid: TAG
transport TAG from diet → tissues

CHM remnants
B-48
E
source: CHM (chylomicrons)
principal lipids: TAG, cholesterol, phospholipids
transport remnants of chylomicrons to liver

VLDL
C-II
B-100
source: liver
principal lipids: TAG
transport newly synthesized TAG to other tissues

IDL
B-100
source: VLDL
principal lipids: cholesterol, phospholipids, TAG
transport VLDL remnants to other tissues

LDL
B-100
source: VLDL
principal lipids: cholesterol
transport cholesterol to extrahepatic tissues

HDL:
A-I
F-II
source: liver
main lipids: cholesterol, lipoproteins
apoproteins supply
transport cholesterol from tissues

back to liver

Type	Source	Principal Lipids	Important apoproteins	they transport:
chylo-microns	intestine	TAG	B-48 C-II E	TAG from diet to tissues
CHM remnants	CHM (chylomicron)	TAG Phospholipid Cholesterol	B-48 E	CHM remnants to liver
VLDL	liver	TAG	B-100, C-II	newly synthesized TAG to other tissues
IDL	VLDL	TAG phospholipids cholesterol	B-100	VLDL remnants to other tissues
LDL	VLDL	cholesterol	B-100	cholesterol to extrahepatic tissues
HDL	liver	cholesterol phospholipids Apoproteins supply	A-I C-II E	cholesterol from tissues back to liver

HDL involved in cholesterol esterification
enzyme LCAT

HDL gets TAG from VLDL
VLDL gets cholesterol ester from HDL
} CETP